

## Pain and Discomfort

**Many people with Fragile X syndrome are not able to communicate that they are experiencing pain and discomfort. Consequently, they may find other ways to let you know that they are in pain.**

---

There is some evidence that challenging behaviour (particularly, [self-injurious behaviour](#)) can occur in response to painful [medical conditions and discomfort](#). However, this does not mean that other forms of challenging behaviour are not influenced by pain and discomfort. For any sudden outbursts of any form of challenging behaviour that appear 'out of the blue' pain and discomfort should be taken into consideration when trying to identify a cause.

Examples of [medical conditions](#) which have been found to lead to self-injurious behaviour in children and adults who have intellectual disability include otitis media (middle ear infections), skin infections and irritation, and gastrointestinal reflux.

Self-injurious behaviour is sometimes directed towards specific areas of the body which might indicate that the person is experiencing pain in that area e.g., banging or rubbing around the ear in response to earache or banging around the jaw in response to toothache. You can see an example of pain related behaviour in the following re-enactment by a role play actor.

;

At other times, the self-injury may be directed towards another part of the body that does not appear to be related. For example children have been observed who pull their own hair in response to chronic stomach pain. Signs of pain are not always obvious.

### The role of peripheral sensory neuropathy

There is some evidence that individuals who have Fragile X syndrome might have a peripheral sensory neuropathy. This disorder can have two important effects that might be relevant to self-injury. First, pain may not be experienced in the 'normal' way. Painful stimuli may not be experienced as painful. Secondly, people may experience unusual sensations in their hands and arms. These sensations are described by some people as pins and needles (dysaesthesia) or a mild burning sensation. If this is the case, then individuals may respond to this sensation by scratching, picking or biting the area in which the tingling is occurring. When this happens in the absence of any pain, this behaviour may become more severe than it would if the

person was able to identify the pain sensation.

[Download: Example 1 - Challenging behaviour and pain](#)

[Download: Example 2 - Challenging behaviour and pain](#)

### **Could self-injurious behaviour relieve pain and discomfort?**

It seems odd that someone would engage in self-injurious behaviour if they were already experiencing pain elsewhere. Pain gate theory suggests that an acute pain (sharp and intense pain such as that caused by self-injurious behaviour) can block (or gate) the sensation of chronic pain (long lasting pain such as that caused by a painful medical condition). Pain gate theory applies not only to individuals who have an intellectual disability but to everyone. Have you ever banged your head and rubbed it better? This is an example of how you are able to block one type of sensation (pain) by creating another.

For information about detecting pain in children with severe intellectual disabilities and/or communication difficulties please [click here](#).

**[NEXT: Sensory stimulation and reinforcement](#)**